

**Original Article**  
**Radiology**

# ASSESSMENT AND EVALUATION OF PATELLOFEMORAL INSTABILITY ON MAGNETIC RESONANCE IMAGING

Sandeep S<sup>1</sup>, Reddy Ravikanth<sup>2</sup>, Babu Philip<sup>3</sup>

<sup>1</sup> - Assistant Professor in Radiology, St. John's Medical College, Bangalore.

<sup>2</sup> - Post-graduate student in Radiology, St. John's Medical College, Bangalore.

<sup>3</sup> - Professor & Head, Department of Radiology, St. John's Medical College, Bangalore.

**Corresponding Author:**

Reddy Ravikanth, MBBS  
Post-graduate, Department of Radiology,  
St. John's Medical College,  
Bangalore - 560034.  
Email: ravikanthreddy06@gmail.com  
Mob: +91-9035144252

Submitted on: 10 June 2016

Accepted on: 21 June 2016

**Abstract:**

**Background:** Patellofemoral instability is a common clinical problem. Risk factors predisposing to patellar dislocation include ligamentous laxity, femoral or tibial malrotation, patella alta or femoral trochlear dysplasia. Objective of this study was to determine the contribution of magnetic resonance imaging (MRI) in evaluating patellofemoral instability.

**Materials and Methods:** 39 patients (45 knees) with patellar instability who underwent knee magnetic resonance imaging between June 2014 and June 2016, at St. John's Medical College Hospital, Bangalore were included. MRI were analyzed for the presence of bone, hyaline cartilage and soft-tissue abnormalities, as well as anatomic variants that may contribute to chronic patellar instability.

**Results:** Of the 45 knees analyzed, we found bone changes in 44%, cartilaginous injuries in 64%, disruption of the medial patellofemoral ligament (MPFL) in 29% and joint morphology abnormalities in 73% patients. Meniscal tears were also identified in 2 (4%) patients and fibular fracture in one (2%) patient.

**Conclusion:** MRI helped in the detection of predisposing factors that may contribute to the development of patellofemoral instability and the diagnosis of bone, hyaline cartilage, ligamentous and meniscal abnormalities.

**Key-words:** patellofemoral joint, instability, MRI

## Introduction:

The patellofemoral joint (PFJ) is a complex structure with high functional and biomechanical requirements. The normal function of this joint is dependent on the congruent relationship of the patella with the trochlear groove. Developmental or acquired alterations in the surface geometry of the PFJ are associated with multiple clinical problems such as patellar instability; chondromalacia patella; and, consequently, anterior knee pain. Therefore, having knowledge about the exact anatomy and function of the joint is required to understand, diagnose, and treat these different abnormalities appropriately. Patellofemoral instability is still a challenge for orthopedists not only because of the diversity and complexity of the causal factors but also because of the large number of therapeutic possibilities. Many authors have taken the view that, in the majority of cases, there is a predisposing anatomical factor that contributes towards the instability and that it is fundamental to recognize this in order to define the best approach.<sup>1</sup> Moreover, displacement of the patella promotes occurrences of other lesions and greater risk of development of osteoarthritis, which also affects the choice of treatment and the prognosis. For many decades, simple radiographs (X-rays) and computed tomography (CT) were the only imaging methods used for evaluating these patients. Magnetic resonance imaging (MRI) has emerged as an auxiliary method for investigating patellofemoral instability, particularly for detecting osteochondral lesions and for evaluating the medial patellofemoral ligament (MPFL).<sup>2</sup> Furthermore, many authors have perceived that MRI also makes it possible to detect predisposing factors

and to make a variety of measurements with the same accuracy as CT scans, thus making this the preferred imaging method for evaluating patellofemoral instability in several centers. The aim of this study was to determine the contribution of MRI in evaluations on patients with a clinical condition of patellofemoral instability.

## Materials and methods:

This study was approved by the Institutional Ethical Review Board (IERB) at St. John's Medical College Hospital, Bangalore. MRI scans on 45 consecutive knee cases were retrospectively analyzed. The patients were referred by orthopedists who are specialists in knee surgery, with a clinical condition of patellofemoral instability and a history of recurrent lateral dislocation of the patella. The MRI scans were produced between June 2014 and June 2016, in apparatus of 1.5 Tesla, using specific coils for the knee joint. All the examinations included T2-weighted sequences with fat suppression in the axial plane, proton density (PD) weighted sequences with fat suppression in the coronal and sagittal planes, and T2-weighted sequences in the sagittal plane. Twenty-three examinations included additional sequences with T1 weighting in the axial and sagittal planes at flexion of around 20° to 30°.

Areas of bone edema typical of previous lateral dislocation of the patella were sought. These were characterized by increased signal in the T2 and PD weightings with fat suppression, in the bone marrow of the medial portion of the patella and on the external face of the lateral femoral condyle, other areas of bone edema unrelated to patellar dislocation, joint effusion and tendinous, ligamentous, meniscal or cartilaginous lesions. The

presence of morphological or joint geometrical abnormalities that would predispose towards patellofemoral instability was also evaluated, using the following criteria:

a) High patella: Caton-Deschamps index greater than 1.2

b) Morphological abnormalities in the intercondylar fossa of the femur (trochlea dysplasia), based on the Dejour radiographic classification:

Type A – trochlea with morphology preserved, but with a shallow trochlear sulcus (angle greater than 145°).

Type B – rectified trochlea.

Type C – asymmetrical trochlear facets, with hypoplasia of the medial facet and convexity of the lateral facet.

Type D – asymmetry of the trochlear facets, with presence of a supra-trochlear ventral prominence greater than or equal to 7 mm.

c) Lateral inclination of the patella: normally, the lateral facet of the patella forms an angle with the posterior bicondylar line that is open laterally and greater than 8° in flexion (lateral inclination angle). Patellae with a lateral inclination angle that opened laterally but was less than or equal to 8° on axial images obtained in flexion were considered to be inclined. If the only images available were axial, in extension, the patellae were considered to be inclined if their inclination angle opened medially or if the lateral facet of the patella was parallel to the posterior bicondylar line.

d) Lateral displacement of the patella: this was only evaluated in the 23 cases in which images acquired in flexion in the axial plane were available, since lateralization of the patella in extension may be physiological. Patellae that did not fit into the trochlea in flexion and with the medial margin lateralized in relation to a line perpendicular to the bicondylar

line, in the plane of the medial eminence of the femoral trochlea, were considered to be displaced.

e) Increased TA-GT distance: this measurement was evaluated only in the cases in which the image acquisition in the axial plane included the insertion of the patellar ligament into the tuberosity of the tibia. Firstly, a line was traced out tangentially to the posterior femoral condyles (posterior bicondylar line). Then, two lines were traced out perpendicular to the posterior bicondylar line: one going through the center of the base of the trochlea (line GT), in the plane in which the intercondylar region has the shape of a Roman arch; and the other going through the center of the patellar ligament (line TA), in the plane of its uppermost insertion into the tuberosity of the tibia, in the first slice in which no fat was identified between the patellar ligament and the cortical bone of the tibia. The distance between the lines TA and GT (TA-GT distance) was measured in millimeters, and this was considered to be within normal limits when it was less than or equal to 15 mm.

### Results:

MRI scans from 45 knees (n = 26 (58%) on the right side and n = 19 (42%) on the left side) were analyzed. These were from 39 patients (six presented bilateral clinical conditions and underwent MRI on both knees), of whom 19 were men (42%) and 26 were women (58%).

The ages of the patients studied ranged from 14 to 53 years, with a mean of 29 years ( $\pm 10$ ) and median of 28 years. Twenty patients (44%) presented a bone contusion pattern typical of recent lateral transitory dislocation of the patella, in association with joint effusion. One

patient presented bone edema relating to an impacted fracture in the head of the fibula, along with signs of recent transitory dislocation of the patella.

Signs of lesions of the patellofemoral ligament were identified in 13 patients (29%), and all of these patients also presented bone edema characteristic of recent dislocation of the patella, with presence of joint effusion and fluid infiltration adjacent to the medial retinaculum.

Thirty-six patients (64%) presented chondral lesions; patella cartilage lesions were identified in 29 patients (51%); and in seven cases (15%), a lesion in the femoral cartilage was also observed. A detached osteochondral fragment could be identified in four of the patients with patella cartilage lesions. Edema of the subchondral bone relating to chondropathy was identified in seven patients (15%), while five patients (11%) presented signs of patellofemoral arthrosis, with reduction of the lateral patellofemoral joint space and presence of osteophytes. The ages of these five patients ranged from 29 to 53 years (mean of  $44 \pm 10$  years). None of these patients presented any signs of arthrosis in the femorotibial compartment.

Morphological or joint geometry abnormalities that predisposed towards patellofemoral instability were found in 33 cases (73%), and these were characterized as follows: high patella (n = 24; 53%); lateral inclination of the patella beyond the physiological limits (n = 25; 56%); patella displaced laterally (n = 13; 29%); and trochlear dysplasia (n = 23; 51%).

An increased TA-GT distance was found in 11 cases (24%), although this measurement could only be made in 26 knees (58% of the cases), because of non-inclusion of the insertion of the patella ligament in the tibial tuberosity,

in the axial plane, in 19 cases. Seventeen patients (38%) presented edema in the superolateral portion of the infrapatellar fat.

**Table 1: Prevalence of findings from MRI**

Abnormalities detected on MRI	n	%
Bone contusion typical of transitory dislocation of the patella	20	44
Joint effusion	22	49
Lesion of the medial patellofemoral ligament	13	29
Chondral lesion:	29	64
Patellar cartilage	29	51
Patellar and femoral cartilages	7	15
Detached osteochondral fragment	4	9
Signal abnormality in the subchondral bone of the patella	7	15
Patellofemoral arthrosis	5	12
Morphological or joint geometry abnormalities:	33	73
High patella	24	53
Laterally inclined patella	25	56
Laterally displaced patella	13	29
Trochlear dysplasia	23	51
Increased TA-GT distance	11	24
Lesion of the medial meniscus	2	4
Fracture of the head of the fibula	1	2

### Discussion:

Patellofemoral instability can be classified as acute or chronic. Acute instability is characterized by traumatic lateral dislocation of the patella and mainly affects adolescents and young adults. Up to 44% of these patients evolve to chronic instability. Chronic instability comprises episodes of recurrent dislocation, with greater risk of development of osteoarthritis.

In the present study, only patients with a history of chronic instability were selected. Their mean age ( $28 \pm 10$  years) was similar to what has been reported in the literature, with predominance of female patients (58%). In the present study group, 44% presented a bone contusion pattern typical of recent dislocation of the patella.

Many authors have taken the view that in most cases of chronic instability, there is a predisposing anatomical factor, and that the more intensely that these factors are present, the lower the intensity of the trauma required for dislocation to occur will be. The commonest predisposing factors are trochlea dysplasia, high patella and lateralization of the anterior tuberosity of the tibia, and recognition of these bone abnormalities is fundamental for defining the prognosis and for achieving the best therapeutic management.<sup>3</sup>

For a long time, radiographic studies on the knee in anteroposterior, absolute lateral and patellar axial views were the main imaging methods available for evaluating the femoropatellar joint. These methods were fundamental for developing concepts relating to the dynamics of the extensor mechanism and for detecting morphological abnormalities of the trochlea.

However, radiographic studies performed in isolation present some important limitations. It is not very easy to obtain images in absolute lateral view, particularly without using fluoroscopy. The lateral view needs to be acquired rigorously, in order to adequately analyze the trochlea, given that rotations of just 5% may generate false positive or false negatives regarding trochlear dysplasia. Lateralization of the tibial

tuberosity has also been implicated in instability, and measurement of the TA-GT distance has become critical data in making therapeutic decisions, since this enables quantitative analysis that is more reliable than the clinical examination.<sup>4</sup>

However, this measurement is also not accurate when made by means of radiographic images, as was demonstrated in the study by Wagenaar et al.<sup>5</sup> For these reasons, CT scans have often been used in association with radiographic studies in evaluating these patients, since this enables measurement of the TA-GT distance, along with assessment of the positioning of the patella and the morphology of the trochlea.<sup>6</sup>

Over the last few years, MRI has also emerged as an auxiliary method for investigating patellofemoral instability, particularly for detecting osteochondral lesions and for evaluating the MPFL, given that X-rays and CT scans are used primarily for bone assessments.<sup>7</sup> Cartilage lesions of the patellofemoral compartment are generally related to instability, as also observed in the present study, in which chondral lesions were found in 64% of the cases. MRI presents sensitivity of around 90% for detecting osteochondral lesions in the knee, which may go unnoticed on X-rays in up to 60% of the cases, thus justifying performing MRI among these patients. Lesions of the joint cartilage predispose towards early arthrosis, and this was observed in five patients (12%), with a mean age of  $44 \pm 10$  years.

MRI also presents excellent results for detecting lesions of the MPFL after acute dislocation of the patella, with estimated sensitivity of up to around 85%. In the present study, lesions of the MPFL were detected in 29% of the

cases, although it has to be emphasized that in all of these cases, there had been a recent episode of dislocation of the patella, with presence of joint effusion and fluid adjacent to the stabilizing structures of the medial compartment, which seems to facilitate detection of fiber discontinuity.<sup>8</sup> Further studies will be needed in order to evaluate the accuracy of MRI for identifying lesions of the MPFL in cases in which there is no recent dislocation.

With greater use of MRI among patients presenting instability, several authors have noticed over recent years that MRI not only has good accuracy in detecting lesions of the cartilage and soft tissues, but also enables detection of predisposing factors.<sup>9</sup> In fact, in the present study, anatomical or joint geometry abnormalities that were considered to be predisposing factors for patellofemoral instability were identified in 73% of the cases. Trochlear dysplasia, which is considered to be one of the main risk factors for instability, was present in more than half of the cases and was easily identified using MRI, thus confirming the results from several authors who had stated that MRI was an excellent method for diagnosing trochlear dysplasia.<sup>10</sup> It was also possible to identify high patella and patellar inclination beyond the physiological limits in 53% and 56% of the cases, respectively, thus confirming the high prevalence of morphological abnormalities of the trochlea and of patellar positioning among these patients.

Measurement of the TA-GT distance is one of the main reasons for requesting tomographic evaluations among patients with patellofemoral instability. With the growing need for evaluation of ligament and joint cartilage integrity, many patients

have started to undergo three types of imaging diagnostics: X-ray, CT scan and MRI, which increases measurements made using CT, including measurement of the TA-GT distance, for MRI, with excellent results. Although comparison between the findings from MRI and CT scans was not among the objectives of the present study, we can affirm that it was possible in the majority of the cases to identify the main morphological and joint geometry abnormalities that predispose towards instability.

In the present study, in which it was not possible to measure the TA-GT distance in 42% of the cases, because of failure to identify the insertion of the patellar ligament on images in the axial plane. One attenuating factor was the fact that a large portion of these patients had already undergone CT examination previously, with tomographic measurement of the TA-GT distance, and MRI was only indicated for detection of other abnormalities. Nonetheless, this indicates that unless special attention is given to inclusion of the tuberosity of the tibia in the axial plane, it can easily be excluded from the study. Another limiting factor of MRI is that because of the physiological lateralization and inclination of the patella in extension, evaluation of these parameters should be done with the knee flexed, which is not part of the habitual routine of MRI on the knee and is not possible on any equipment. Despite these limitations, the capacity to combine evaluation on bone, cartilage and soft-tissue structures has meant that in many centers, MRI is today considered to be the preferred imaging method for evaluating patellofemoral instability, thus replacing CT, with the additional benefit of not exposing patients to the risks of ionizing radiation.

### Conclusion:

MRI serves as an excellent imaging modality in making the diagnosis, detecting factors that predispose towards patellofemoral instability and identifying lesions that tend not to be identified on X-rays and CT scans, such as cartilaginous lesions and those of the MPFL. Evaluation of trochlear measurements is also important when evaluating MR images in patients with patellofemoral instability.

Conflict of interest: None

Source of funding: None

Informed consent: Obtained

Ethical clearance: Taken from Institutional Ethical Review Board, St. John's Medical College, Bangalore – 560 034.

### References:

1. Dejour H, Walch G, Nove-Josserand L, Guier C. Factors of patellar instability: an anatomic radiographic study. *Knee Surg Sports Traumatol Arthrosc* 1994; 2:19–26.
2. Christoforakis JJ, Strachan RK. Internal derangements of the knee associated with patellofemoral joint degeneration. *Knee Surg Sports Traumatol Arthrosc* 2005; 13:581–584.
3. Stäubli HU, Dürrenmatt U, Porcellini B, Rauschnin W. Anatomy and surface geometry of the patellofemoral joint in the axial plane. *J Bone Joint Surg Br* 1999; 81:452–45.
4. Kijowski R, Blankenbaker DG, Davis KW, Shinki K, Kaplan LD, De Smet AA. Comparison of 1.5-T and 3.0-T MR imaging for evaluating the articular cartilage of the knee joint. *Radiology* 2009; 250:839–848.
5. Biedert RM, Bachmann M. Anterior – posterior

trochlear measurements of normal and dysplastic trochlea by axial magnetic resonance imaging. *Knee Surg Sports Traumatol Arthrosc*. 2009;17(10):1225-30.

6. Christian SR, Anderson MB, Workman R, Conway WF, Pope TL. Imaging of anterior knee pain. *Clin Sports Med*. 2006;25(4):681-702.
7. Merchant AC. Patellofemoral Imaging. *Clin Orthop Relat Res*. 2001;(389):15-21.
8. Keser S, Savranlar A, Bayar A, Ege A, Turhan E. Is there a relationship between anterior knee pain and femoral trochlear dysplasia? Assessment of lateral trochlear inclination by magnetic resonance imaging. *Knee Surg Sports Traumatol Arthrosc*. 2008;16(10):911-5.
9. Lim AKS, Chang HC, Hui JHP. Recurrent patellar dislocation: reappraising our approach to surgery. *Ann Acad Med Singapore*. 2008;37(4):320-3.
10. Schoettle PB, Zanetti M, Seifert B, Pfirrmann CWA, Fucentese SF, Romero J. The tibial tuberosity–trochlear groove distance; a comparative study between CT and MRI scanning. *The Knee*. 2006;13(1):26-31.